Problem A1

Let A be a positive real number. What are the possible values of $\sum_{j=0}^{\infty} x_j^2$, given that $x_0, x_1, x_2, \ldots$ are positive numbers for which $\sum_{j=0}^{\infty} x_j = A$?

Problem A2

Prove that there exist infinitely many integers n such that n, $n+1$, and $n+2$ are each the sum of two squares of integers.
[*Example:* $0 = 0^2 + 0^2$, $1 = 0^2 + 1^2$, and $2 = 1^2 + 1^2$.]

Problem A3

The octagon $P_1P_2P_3P_4P_5P_6P_7P_8$ is inscribed in a circle, with the vertices around the circumference in the given order. Given that the polygon $P_1P_3P_5P_7$ is a square of area 5 and the polygon $P_2P_4P_6P_8$ is a rectangle of area 4, find the maximum possible area of the octagon.

Problem A4

Show that the improper integral

$$\lim_{B\to\infty} \int_0^B \sin(x)\, \sin(x^2)\, dx$$

converges.

Problem A5

Three distinct points with integer coordinates lie in the plane on a circle of radius $r > 0$. Show that two of these points are separated by a distance of at least $r^{1/3}$.

Problem A6

Let $f(x)$ be a polynomial with integer coefficients. Define a sequence $a_0, a_1, \ldots$ of integers such that $a_0 = 0$ and $a_{n+1} = f(a_n)$ for all $n \geq 0$. Prove that if there exists a positive integer m for which $a_m = 0$ then either $a_1 = 0$ or $a_2 = 0$.

Problem B1

Let a_j, b_j, and c_j be integers for $1 \leq j \leq N$. Assume, for each j, that at least one of a_j, b_j, c_j is odd. Show that there exist integers r, s, t such that $ra_j + sb_j + tc_j$ is odd for at least $4N/7$ values of j, $1 \leq j \leq N$.

Problem B2

Prove that the expression

$$\frac{\gcd(m,n)}{n}\binom{n}{m}$$

is an integer for all pairs of integers $n \geq m \geq 1$. [Here $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ and $\gcd(m,n)$ is the greatest common divisor of m and n.]

Problem B3

Let $f(t) = \sum_{j=1}^{N} a_j \sin(2\pi jt)$, where each a_j is real and $a_N \neq 0$. Let N_k denote the number of zeros (including multiplicities) of $\frac{d^k f}{dt^k}$. Prove that

$$N_0 \leq N_1 \leq N_2 \leq \dots \qquad \text{and} \lim_{k\to\infty} N_k = 2N.$$

Problem B4

Let $f(x)$ be a continuous function such that $f(2x^2-1) = 2\,x\,f(x)$ for all x. Show that $f(x) = 0$ for $-1 \leq x \leq 1$.

Problem B5

Let S_0 be a finite set of positive integers. We define finite sets $S_1, S_2, \dots$ of positive integers as follows:

Integer a is in S_{n+1} if and only if exactly one of $a-1$ or a is in S_n.

Show that there exist infinitely many integers N for which $S_N = S_0 \cup \{N + a : a \in S_0\}$.

Problem B6

Let B be a set of more than $2^{n+1}/n$ distinct points with coordinates of the form $(\pm 1, \pm 1, \dots, \pm 1)$ in n-dimensional space, with $n \geq 3$. Show that there are three distinct points in B which are the vertices of an equilateral triangle.