

1. FORMULAS FOR EXPONENTIAL AND POWER EXPRESSIONS

$1^a = 1$ for any a , and $a^0 = 1$ if $a \neq 0$

$$x^{a+b} = x^a \cdot x^b$$

$$x^{a-b} = \frac{x^a}{x^b}$$

$$x^{a \cdot b} = (x^a)^b$$

$$x^a \cdot y^a = (x \cdot y)^a$$

$$\frac{x^a}{y^a} = \left(\frac{x}{y}\right)^a$$

2. FORMULAS FOR RADICAL EXPRESSIONS

$\sqrt[n]{1} = 1$ for any $n \geq 2$, and $\sqrt[n]{0} = 0$ for any $n \geq 2$

$\sqrt[n]{-1}$ undefined if n even, and $\sqrt[n]{-1} = -1$ if n odd

$$\sqrt[n]{x} \cdot \sqrt[n]{y} = \sqrt[n]{x \cdot y}$$

$$\frac{\sqrt[n]{x}}{\sqrt[n]{y}} = \sqrt[n]{\frac{x}{y}}, \text{ if } y \neq 0$$

3. RADICAL EXPRESSIONS CONVERTED TO EXPONENTIAL EXPRESSIONS

$$\sqrt[n]{x} = x^{1/n}$$

$$\sqrt[n]{x^n} = \begin{cases} x, & \text{if } n \text{ is odd;} \\ |x|, & \text{if } n \text{ is even.} \end{cases}$$

$$(\sqrt[n]{x})^n = x$$

$$\sqrt[n]{x^m} = (\sqrt[n]{x})^m = x^{m/n}$$

4. NEGATIVE POWERS FOR EXPONENTIAL AND RADICAL EXPRESSIONS

$$x^{-n} = \frac{1}{x^n}$$

$$\left(\frac{x}{y}\right)^{-n} = \left(\frac{y}{x}\right)^n$$

$$x^{-1/n} = \frac{1}{\sqrt[n]{x}}$$

$$x^{-m/n} = \frac{1}{\sqrt[n]{x^m}} = \frac{1}{(\sqrt[n]{x})^m}$$

$$\left(\frac{x}{y}\right)^{-1/n} = \sqrt[n]{\frac{y}{x}}$$

5. FORMULAS FOR FACTORING SPECIAL POLYNOMIALS

5.1. Difference of two squares.

$$x^2 - a^2 = (x - a) \cdot (x + a)$$

5.2. Sum of two squares.

$x^2 + a^2$ is prime, cannot be factored

5.3. Difference of two cubes.

$$x^3 - a^3 = (x - a) \cdot (x^2 + a \cdot x + a^2)$$

5.4. Sum of two cubes.

$$x^3 + a^3 = (x + a) \cdot (x^2 - a \cdot x + a^2)$$

5.5. Zero-factor Property.

If $a \cdot b = 0$, then $a = 0$ or $b = 0$

6. FORMULAS FOR EXPANDING SPECIAL POLYNOMIALS

6.1. The Square of the Sum.

$$(x + a)^2 = x^2 + 2ax + a^2$$

6.2. The Square of the Difference.

$$(x - a)^2 = x^2 - 2ax + a^2$$

6.3. The Cube of the Sum.

$$(x + a)^3 = x^3 + 3ax^2 + 3a^2x + a^3$$

6.4. The Cube of the Difference.

$$(x - a)^3 = x^3 - 3ax^2 + 3a^2x - a^3$$

7. EQUATIONS AND INEQUALITIES WITH ABSOLUTE VALUES

If $x \geq 0$, then $|x| = x$.

If $x < 0$, then $|x| = -x$.

If $k > 0$, then

$$\begin{aligned} |x| = k & \text{ is equivalent to } x = k \text{ or } x = -k \\ |x| < k & \text{ is equivalent to } -k < x < k \\ |x| > k & \text{ is equivalent to } x < -k \text{ or } x > k \end{aligned}$$

8. COMPLEX NUMBERS

8.1. The Definition of i .

$$\sqrt{-1} = i, \quad i^2 = -1$$

If a is strictly positive, then

$$\sqrt{-a} = i \cdot \sqrt{a} = \sqrt{a} i$$

8.2. Sum of Two Complex Numbers.

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

8.3. Difference of Two Complex Numbers.

$$(a + bi) - (c + di) = (a - c) + (b - d)i$$

8.4. Multiplication of Two Complex Numbers.

$$(a + bi) \cdot (c + di) = (ac - bd) + (ad + bc)i$$

8.5. The Length of a Complex Number or its Absolute Value.

$$|a + bi| = \sqrt{a^2 + b^2}$$

8.6. The Conjugate of a Complex Number.

The conjugate of $a + bi$ is $a - bi$

8.7. Multiplying Two Conjugate Complex Numbers.

$$(a + bi) \cdot (a - bi) = a^2 + b^2$$

9. DISTANCE FORMULA BETWEEN TWO POINTS

The distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by the formula

$$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

10. LINES AND SLOPES

The slope of the line passing through the points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is given by the formula

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

The slope of a horizontal line $y = a$ is $m = 0$, while the slope of a vertical line $x = b$ is undefined.

If the slope of the line $y = mx + b$ is strictly positive, the line is strictly increasing. If the slope of the line $y = mx + b$ is strictly negative, the line is strictly decreasing.

The slope-intercept form of a line is

$$y = mx + b \quad ,$$

where m is the slope, and b is the y intercept.

The point-slope form of a line is

$$y - y_1 = m(x - x_1) \quad ,$$

where m is the line's slope, and (x_1, y_1) is a given point on the line.

Two lines with the same slope are parallel. If the product of the slopes of two lines is -1 , the lines are perpendicular.

11. SOLVING QUADRATIC EQUATIONS

The quadratic equations are of the form $ax^2 + bx + c = 0$. If you can factor $ax^2 + bx + c$ as a product of two terms, then you can use the 0-property to solve it. Let see other methods to solve it.

11.1. Solving equations of the form $x^2 = a$.

$$x^2 = a \Rightarrow \sqrt{x^2} = \sqrt{a} \Rightarrow |x| = \sqrt{a} \Rightarrow x = \sqrt{a} \text{ or } x = -\sqrt{a}$$

11.2. **Completing the square.** Consider the equation $x^2 + bx + c = 0$ (the coefficient of x^2 has to be 1). Then, you complete the square:

$$\begin{aligned}
 x^2 + bx + c = 0 &\Rightarrow x^2 + bx = -c \Rightarrow x^2 + bx + \left(\frac{b}{2}\right)^2 = \left(\frac{b}{2}\right)^2 - c \Rightarrow \left(x + \frac{b}{2}\right)^2 = \left(\frac{b}{2}\right)^2 - c \Rightarrow \\
 &\Rightarrow \sqrt{\left(x + \frac{b}{2}\right)^2} = \sqrt{\left(\frac{b}{2}\right)^2 - c} \Rightarrow \left|x + \frac{b}{2}\right| = \sqrt{\left(\frac{b}{2}\right)^2 - c} \Rightarrow x + \frac{b}{2} = \pm \sqrt{\left(\frac{b}{2}\right)^2 - c} \Rightarrow \\
 &\Rightarrow x = -\frac{b}{2} + \sqrt{\left(\frac{b}{2}\right)^2 - c} \quad \text{or} \quad x = -\frac{b}{2} - \sqrt{\left(\frac{b}{2}\right)^2 - c}
 \end{aligned}$$

11.3. **Quadratic formula.** The solutions of the quadratic equation $ax^2 + bx + c = 0$ are :

$$x_1 = \frac{-b + \sqrt{b^2 - 4 \cdot a \cdot c}}{2 \cdot a}, \quad x_2 = \frac{-b - \sqrt{b^2 - 4 \cdot a \cdot c}}{2 \cdot a}$$

The discriminant $\Delta = b^2 - 4ac$:

If	then the roots of
$b^2 - 4ac > 0$	$ax^2 + bx + c$ are
$b^2 - 4ac = 0$	real and unequal
$b^2 - 4ac < 0$	real and equal
	complex and conjugate